

**KELIPS® Gel***Protective covering gel***INDICATION:**

**KELIPS® GEL** is indicated for the symptomatic treatment of cheilitis resulting from an isotretinoin-based acne treatment.

**COMPOSITION OF KELIPS® Gel:**

Oxidised triglycerides (OGT) from plant-based essential fatty acids (87.50%).  
Excipients: silicon dioxide (7.00%), zinc oxide (1.00%), perfume (4.00%).

**PRESENTATION:**

**KELIPS® Gel:** 15G tube

**CHEILITIS? ORIGIN AND SYMPTOMS**

**Cheilitis** can be caused by infections or medication. Cases of **isotretinoin-based acne treatments** present a 100% rate of patients with cheilitis.

**Cheilitis** symptoms include dry or very dry lips with permanent burning sensations and painful cracking. They can make speaking and eating difficult due to limited and painful opening of the mouth or abnormal sticking of the upper and lower lip during the night and painful separation of the two lips and the aggravation of existing cracks when the lips start moving in the morning. **Cheilitis affect the quality of life for patients treated by isotretinoin.**

**KELIPS® gel MODE OF ACTION**

Thanks to its invisible protective film made of OGT (oxidised triglycerides made from plant-based essential fatty acids), **KELIPS® GEL covers the cheilitis and protects them against external aggression** (Cheilitis symptoms: dry/irritated/red/erythematous/cracked lips). **KELIPS® GEL** consists of 87.5% OGT. The OGT have a bioactive action on the tissues (Cheilitis zone) and OGT promote the natural process of healing/regeneration. In this way, **KELIPS® GEL** helps the cheilitis healing and scarring process.

**EFFICIENCY OF KELIPS® Gel**

Efficiency of **KELIPS® Gel** was demonstrated in a clinical study\*.

**Used as a preventive treatment**, the OGT gel proved efficient in 27% of the patients who fully observed their **KELIPS® GEL** treatment.

**Used as a curative treatment**, **KELIPS® Gel** significantly reduced all signs and symptoms of cheilitis and the discomfort they cause due to their intensity in most of the study's patients.

**The benefits brought by KELIPS® Gel** were all the more important when the patients observed the **KELIPS® Gel** treatment.

\* *GOAL study evaluating the preventive and curative properties of **KELIPS® Gel** in patients presenting cheilitis resulting from an isotretinoin-based acne treatment.*

**INSTRUCTIONS FOR USE:**

Apply **KELIPS® Gel** on the lips **three times a day**, and more if necessary, preferably after meals and for the entire duration of the acne treatment.

KELIPS® Gel must be stored at room temperature and out of children's reach.

Do not use after the expiry date stamped on the crimp of the tube and on the case.

A 15g tube is enough of a one-month treatment.

**WARNING:**

**KELIPS® Gel** is for external use only and should not be used in the eye area.

The product's harmlessness and good tolerance have been demonstrated.

**KELIPS® Gel** should not be used in the case of an over-infected lesion as well as on people with hypersensitivity to one of the components. In the event of an allergic reaction, we recommend interrupting the treatment and seeking medical advice.

Do not use on children under the age of **12**. Do not use in pregnant or lactating women.

Do not use if the packaging is damaged.

**KELIPS® Gel** should not be used in the same area in conjunction with other products in without medical advice.

MEDICAL DEVICE



---

**Year of CE marking:** XXXX

**Last revised:** XXXX



**Manufacturer:** Laboratoires Carilene, 7, rue du Chant des Oiseaux - 78360 Montesson – France.

**Distributor:** xxxxxxxxxxxx



“Please read the instructions for use”



2007/47/EC”

“Medical device compliant with the essential requirements of Directive 93/42/EE and



“To be used until”



“Batch number”



“Manufacturer”